

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 5/26/99	3. REPORT TYPE AND DATES COVERED Final Report 9/1/93 - 8/31/96		
4. TITLE AND SUBTITLE Cloning of P8, A Transcription Factor Required of Skeletogenic Gene Expression		5. FUNDING NUMBERS N00014-93-1-1400		
6. AUTHOR(S) PI: Eric H. Davidson		8. PERFORMING ORGANIZATION REPORT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Division of Biology California Institute of Technology 1200 E. California Blvd. Pasadena, CA 91125		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5000		11. SUPPLEMENTARY NOTES <div style="font-size: 2em; font-weight: bold; text-align: center;">19990601 082</div>		
12a. DISTRIBUTION/AVAILABILITY STATEMENT Distribution Unlimited		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) The transcription factor toward which this project was directed (formerly known as P8, now as Sp(G/C)F-1) interacts with regulatory regions of many sea urchin embryo genes, and probably acts as a general enhancer protein. During this year this factor was: <i>i</i> , purified by affinity chromatography; <i>ii</i> , microsequence was obtained; <i>iii</i> , the factor was sequenced; <i>iv</i> , its DNA binding site was determined; <i>v</i> , it was expressed in bacteria and an antibody generated; <i>vi</i> , its provenance was determined; and an unusual aspect of its structure was analyzed. The latter is that the factor appears in sea urchin embryo nuclear extract in five different forms of decreasing molecular weight. These forms are also presented in cell-free translation using synthetic mRNA from the cloned cDNA. We showed that—unusually—there are five successive ATG start sites in this mRNA, all used more or less equally, thus accounting for five nested forms. The sequence reveals Sp(G/C)F-1 to be a novel DNA-binding protein, unlike any in the data banks. It is loaded into the egg in relatively large quantities during oogenesis and then appears in embryo nuclei.				
14. SUBJECT TERMS sea urchin, transcription factor, embryo		15. NUMBER OF PAGES 16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

FINAL REPORT

Grant #: N00014-93-1-1400

PRINCIPAL INVESTIGATOR: Eric H. Davidson

INSTITUTION: California Institute of Technology

GRANT TITLE: Cloning of P8, A Transcription Factor Required for Skeletogenic Gene Expression

AWARD PERIOD: 9/1/93 - 8/31/96

OBJECTIVE: This is an AASERT Graduate Training Project, the object of which is to isolate, purify, clone, and characterize transcription factors that interact with the regulatory domains of skeletogenic genes in the sea urchin.

APPROACH: The known target site of the factor Sp(G/C)F-1, which interacts with the SM50 and SM30 skeletogenic genes, provides the means to purify this protein, clone it and characterize its interaction with DNA, its own structure, its provenance, and its particular function.

ACCOMPLISHMENTS: The transcription factor toward which this project was directed (formerly known as P8, now as Sp(G/C)F-1) interacts with regulatory regions of many sea urchin embryo genes, and probably acts as a general enhancer protein. During this year this factor was: *i*, purified by affinity chromatography; *ii*, microsequence was obtained; *iii*, the factor was sequenced; *iv*, its DNA binding site was determined; *v*, it was expressed in bacteria and an antibody generated; *vi*, its provenance was determined; and an unusual aspect of its structure was analyzed. The latter is that the factor appears in sea urchin embryo nuclear extract in five different forms of decreasing molecular weight. These forms are also presented in cell-free translation using synthetic mRNA from the cloned cDNA. We showed that—unusually—there are five successive ATG start sites in this mRNA, all used more or less equally, thus accounting for five nested forms. The sequence reveals Sp(G/C)F-1 to be a novel DNA-binding protein, unlike any in the data banks. It is loaded into the egg in relatively large quantities during oogenesis and then appears in embryo nuclei.

SIGNIFICANCE: In the course of this project the Graduate Fellow carrying out the work acquired professional expertise in the complete range of molecular technologies required for transcription factor isolation and characterization. As a result we now have our hands on another of the regulatory molecules that control gene expression, including skeletogenic gene expression.

PATENT INFORMATION: None

AWARD INFORMATION: Received Ph.D.

PUBLICATIONS:

Zeller, R. W., Coffman, J. A., Harrington, M. G., Britten, R. J. and Davidson, E. H. SpGCF1, a sea urchin embryo transcription factor, exists as five nested variants encoded by a single mRNA. *Dev. Biol.* **169**, 713-727, 1995.

Zeller, R. W., Britten, R. J. and Davidson, E. H. Developmental utilization of SpP3A1 and SpP3A2: Two proteins which recognize the same DNA target site in several sea urchin gene regulatory regions. *Dev. Biol.* **170**, 75-82, 1995.

Zeller, R. W., Griffith, J. D., Moore, J. G., Kirchhamer, C. V., Britten, R. J. and Davidson, E. H. A multimerizing transcription factor of sea urchin embryos capable of looping DNA. *Proc. Natl. Acad. Sci. USA* **92**, 2989-2993, 1995.